



"Committed to the protection and promotion of public health."

Gallatin City-County Health Department

****Health Alert Network (HAN) Message****

UPDATED INFORMATION ON SCHOOL CLOSURES

Message Type: Update
To: Gallatin Schools
Date: May 7, 2009
Re: School Closure Recommendations

[State of Montana DPHHS HAN UPDATE](#) Distributed via the DPHHS Health Alert Network (HAN) System Wednesday, May 6, 2009 10:20 AM

New information on disease severity and the extent of community spread warrant revision of the CDC's school closure guidance. At this time, CDC recommends the primary means to reduce spread of influenza in schools is to focus on early identification of ill students and staff, staying home when ill, and good cough and hand hygiene etiquette. Decisions about school closure should be at the discretion of local authorities based on local considerations, including public concern and the impact of school absenteeism and staffing shortages.

Montana's Office of Public Instruction (OPI) has distributed information to schools and is emphasizing working closely with local and state health authorities as school issues are identified.

Recommendations include:

- School closure is **not** advised for a suspected or confirmed case of novel influenza A (H1N1) and, in general, is not advised unless there is a magnitude of faculty or student absenteeism that interferes with the school's ability to function.
- Schools that were closed based on previous interim CDC guidance related to this outbreak may reopen.
- Students, faculty or staff with influenza-like illness (fever with a cough or sore throat) should stay home and not attend school or go into the community, except to seek medical care, for at least 7 days even if symptoms resolve sooner.
- Students, faculty and staff who are still sick 7 days after they become ill should continue to stay home from school until at least 24 hours after symptoms have resolved.
- Students, faculty and staff who appear to have an influenza-like illness at arrival or become ill during the school day should be isolated promptly in a room separate from other students and sent home.

- Parents and guardians should monitor their school-aged children, and faculty and staff should self-monitor every morning for symptoms of influenza-like illness.
- Ill students should not attend alternative child care or congregate in settings other than school.
- School administrators should communicate regularly with local public health officials to obtain guidance about reporting of influenza-like illnesses in the school.
- Schools can help serve as a focus for educational activities aimed at promoting ways to reduce the spread of influenza, including hand hygiene and cough etiquette.
- Students, faculty and staff should stringently follow sanitary measures to reduce the spread of influenza, including covering their nose and mouth with a tissue when coughing or sneezing (or coughing or sneezing into their sleeve if a tissue isn't available), frequently washing hands with soap and water, or using hand sanitizer if hand washing with soap and water is not possible.

Categories of Health Alert Messages:

Health Alert	Conveys the highest level of importance; warrants immediate action or attention.
Health Advisory	Provides important information for a specific incident or situation; may not require immediate action.
Health Update	Provides updated information regarding an incident or situation; unlikely to require immediate action.